

Early Imaging AV-133 Imaging

3 of 15 completed

Form Instructions

Women of childbearing potential must have a negative urine and serum pregnancy test result prior to the baseline imaging scan and must have a negative urine pregnancy test result prior to injection of a follow up imaging scan.

Assessment Date

10/12/2020



Vital Signs Measured Approximately 5 Minutes Prior to Injection

1. Was a study physician (or designee) present to evaluate the participant prior to injection? ⋮

No

Yes

If no, please explain: ⋮

Input text

2. Time vital signs measures prior to injection: ⋮

00:00



3a. Supine blood pressure: systolic (mmHg) ⋮

To be taken after participant has been supine for 1-3 minutes)

Input a number

3b. Supine blood pressure: diastolic (mmHg)



To be taken after participant has been supine for 1-3 minutes)

Input a number

4. Supine heart rate: (beats per minute)



To be taken after participant has been supine for 1-3 minutes)

Input a number

5. Time of 18-AV-133 injection:



00:00



Vital Signs Measured Approximately 15 Minutes Post-Injection

6. Time vital signs measured after injection:



00:00



7a. Supine blood pressure: systolic (mmHg)



To be taken after participant has been supine for 1-3 minutes)

Input a number

7b. Supine blood pressure: diastolic (mmHg)



To be taken after participant has been supine for 1-3 minutes)

Input a number

8. Supine heart rate: (beats per minute)



To be taken after participant has been supine for 1-3 minutes)

Input a number

9. AV-133 PET imaging scan:

Not Completed

Completed



If not collected, provide reason:



Input text

10. Was a study physician present to evaluate the participant prior to injection?



No

Yes

If no, please explain:



Input text



Early Imaging Pregnancy Test

Magnetic Resonance Imaging (MRI)

